

Cabbage

Cabbage or **headed cabbage** (comprising several cultivars of *Brassica oleracea*) is a leafy green, red (purple), or white (pale green) biennial plant grown as an annual vegetable crop for its dense-leaved heads. It is descended from the wild cabbage, *B. oleracea* var. *oleracea*, and belongs to the "cole crops", meaning it is closely related to broccoli and cauliflower (var. *botrytis*); Brussels sprouts (var. *gemmifera*); and savoy cabbage (var. *sabauda*). *Brassica rapa* is commonly named Chinese, celery or napa cabbage and has many of the same uses.

Cabbage heads generally range from 0.5 to 4 kilograms (1 to 9 lb), and can be green, purple or white. Smooth-leafed, firm-headed green cabbages are the most common. Smooth-leafed purple cabbages and crinkle-leafed savoy cabbages of both colors are more rare. It is a multi-layered vegetable. Under conditions of long sunny days, such as those found at high northern latitudes in summer, cabbages can grow quite large. As of 2012, the heaviest cabbage was 62.71 kilograms (138.25 lb).

Cabbage was most likely domesticated somewhere in Europe before 1000 BC, although savoys were not developed until the 16th century AD. By the Middle Ages, cabbage had become a prominent part of European cuisine. Cabbage heads are generally picked during the first year of the plant's life cycle, but plants intended for seed are allowed to grow a second year, and must be kept separate from other cole crops to prevent cross-pollination. Cabbage is prone to several nutrient deficiencies, as well as to multiple pests, and bacterial and fungal diseases.

Cabbages are prepared many different ways for eating; they can be pickled, fermented (for dishes such as sauerkraut), steamed, stewed, sautéed, braised, or eaten raw. Cabbage is a good source of vitamin K, vitamin C and dietary fiber. The Food and Agriculture Organization of the United Nations (FAO) reported that world production of cabbage and other brassicas for 2014 was 71.8 million metric tonnes, with China accounting for 47% of the world total.

Cabbage



A whole white cabbage and a longitudinal section

Species	<i>Brassica oleracea</i>
Cultivar group	Capitata Group
Origin	Europe, prior to 1000 BC
Cultivar group members	White cabbage Red cabbage Savoy cabbage

Contents

Taxonomy and etymology

Description

History

Cultivation

Cultivars

Cultivation problems

Production

Culinary use

Nutrients and phytochemicals

Herbal medicine

Disadvantages

Bloating

Food-borne illness

Goiter and iodine intake

See also

References

Works cited

External links

Taxonomy and etymology

Cabbage (*Brassica oleracea* or *B. oleracea* var. *capitata*,^[1] var. *tuba*, var. *sabauda*^[2] or var. *acephala*)^[3] is a member of the genus *Brassica* and the mustard family, Brassicaceae. Several other cruciferous vegetables (sometimes known as *cole crops*^[2]) are considered cultivars of *B. oleracea*, including broccoli, collard greens, brussels sprouts, kohlrabi and sprouting broccoli. All of these developed from the wild cabbage *B. oleracea* var. *oleracea*, also called colewort or field cabbage. This original species evolved over thousands of years into those seen today, as selection resulted in cultivars having different characteristics, such as large heads for cabbage, large leaves for kale and thick stems with flower buds for broccoli.^[1]



Cabbage

The varietal epithet *capitata* is derived from the Latin word for "having a head".^[4] *B. oleracea* and its derivatives have hundreds of common names throughout the world.^[5]

"Cabbage" was originally used to refer to multiple forms of *B. oleracea*, including those with loose or non-existent heads.^[6] A related species, *Brassica rapa*, is commonly named Chinese, napa or celery cabbage, and has many of the same uses.^[7] It is also a part of common names for several unrelated species. These include cabbage bark or cabbage tree (a member of the genus *Andira*) and cabbage palms, which include several genera of palms such as *Mauritia*, *Roystonea oleracea*, *Acrocomia* and *Euterpe oenocarpus*.^{[8][9]}

The original family name of brassicas was *Cruciferae*, which derived from the flower petal pattern thought by medieval Europeans to resemble a crucifix.^[10] The word *brassica* derives from *bresic*, a Celtic word for cabbage.^[6] Many European and Asiatic names for cabbage are derived from the Celto-Slavic root *cap* or *kap*, meaning "head".^[11] The late Middle English word *cabbage* derives from the word *caboche* ("head"), from the Picard dialect of Old French. This in turn is a variant of the Old French *caboce*.^[12] Through the centuries, "cabbage" and its derivatives have been used as slang for numerous items, occupations and activities. Cash and tobacco have both been described by the slang "cabbage", while "cabbage-head" means a fool or stupid person and "cabbaged" means to be exhausted or, vulgarly, in a vegetative state.^[13]

Description

Cabbage seedlings have a thin taproot and cordate (heart-shaped) cotyledon. The first leaves produced are ovate (egg-shaped) with a lobed petiole. Plants are 40–60 cm (16–24 in) tall in their first year at the mature vegetative stage, and 1.5–2.0 m (4.9–6.6 ft) tall when flowering in the second year.^[14] Heads average between 0.5 and 4 kg (1 and 8 lb), with fast-growing, earlier-maturing varieties producing smaller heads.^[15] Most cabbages have thick, alternating leaves, with margins that range from wavy or lobed to highly dissected; some varieties have a waxy bloom on the leaves. Plants have root systems that are fibrous and shallow.^[10] About 90 percent of the root mass is in the upper 20–30 cm (8–12 in) of soil; some lateral roots can penetrate up to 2 m (6.6 ft) deep.^[14]

The inflorescence is an unbranched and indeterminate terminal raceme measuring 50–100 cm (20–40 in) tall,^[14] with flowers that are yellow or white. Each flower has four petals set in a perpendicular pattern, as well as four sepals, six stamens, and a superior ovary that is two-celled and contains a single stigma and style. Two of the six stamens have shorter filaments. The fruit is a silique that opens at maturity through dehiscence to reveal brown or black seeds that are small and round in shape. Self-pollination is impossible, and plants are cross-pollinated by insects.^[10] The initial leaves form a rosette shape comprising 7 to 15 leaves, each measuring 25–35 cm (10–14 in) by 20–30 cm (8–12 in);^[14] after this, leaves with shorter petioles develop and heads form through the leaves cupping inward.^[2]

Many shapes, colors and leaf textures are found in various cultivated varieties of cabbage. Leaf types are generally divided between crinkled-leaf, loose-head savoys and smooth-leaf firm-head cabbages, while the color spectrum includes white and a range of greens and purples. Oblate, round and pointed shapes are found.^[16]

Cabbage has been selectively bred for head weight and morphological characteristics, frost hardiness, fast growth and storage ability. The appearance of the cabbage head has been given importance in selective breeding, with varieties being chosen for shape, color, firmness and other physical characteristics.^[17] Breeding objectives are now focused on increasing resistance to various insects and diseases and improving the nutritional content of cabbage.^[18] Scientific research into the genetic modification of *B. oleracea* crops, including cabbage, has included European Union and United States explorations of greater insect and herbicide resistance.^[19]

History

Although cabbage has an extensive history,^[20] it is difficult to trace its exact origins owing to the many varieties of leafy greens classified as "brassicas".^[21] The wild ancestor of cabbage, *Brassica oleracea*, originally found in Britain and continental Europe, is tolerant of salt but not encroachment by other plants and consequently inhabits rocky cliffs in cool damp coastal habitats,^[22] retaining water and nutrients in its slightly thickened, turgid leaves. According to the triangle of U theory of the evolution and relationships between *Brassica* species, *B. oleracea* and other closely related kale vegetables (cabbages, kale, broccoli, Brussels sprouts, and cauliflower) represent one of three ancestral lines from which all other brassicas originated.^[23]

Cabbage was probably domesticated later in history than Near Eastern crops such as lentils and summer wheat. Because of the wide range of crops developed from the wild *B. oleracea*, multiple broadly contemporaneous domestications of cabbage may have occurred throughout Europe. Nonheading cabbages and kale were probably the first to be domesticated, before 1000 BC,^[24] by the Celts of central and western Europe.^[6]

Unidentified brassicas were part of the highly conservative unchanging Mesopotamian garden repertory.^[25]

It is believed that the ancient Egyptians did not cultivate cabbage,^[26] which is not native to the Nile valley, though a word *shaw't* in Papyrus Harris of the time of Ramesses III, has been interpreted as "cabbage".^[27] Ptolemaic Egyptians knew the cole crops as *gramb*, under the influence of Greek *krambe*, which had been a familiar plant to the Macedonian antecedents of the Ptolemies,^[27] By early Roman times Egyptian artisans and children were eating cabbage and turnips among a wide variety of other vegetables and pulses.^[28]

The ancient Greeks had some varieties of cabbage, as mentioned by Theophrastus, although whether they were more closely related to today's cabbage or to one of the other *Brassica* crops is unknown.^[24] The headed cabbage variety was known to the Greeks as *krambe* and to the Romans as *brassica* or *olus*;^[29] the open, leafy variety (kale) was known in Greek as *raphanos* and in Latin as *caulis*.^[29]

Chrysippus of Cnidos wrote a treatise on cabbage, which Pliny knew,^[30] but it has not survived. The Greeks were convinced that cabbages and grapevines were inimical, and that cabbage planted too near the vine would impart its unwelcome odor to the grapes; this Mediterranean sense of antipathy survives today^[31]



The cabbage inflorescence, which appears in the plant's second year of growth, features white or yellow flowers, each with four perpendicularly arranged petals.



Cabbage with Moong-dal Curry

Brassica was considered by some Romans a table luxury,^[32] although Lucullus considered it unfit for the senatorial table.^[33] The more traditionalist Cato the Elder, espousing a simple, Republican life, ate his cabbage cooked or raw and dressed with vinegar; he said it surpassed all other vegetables, and approvingly distinguished three varieties; he also gave directions for its medicinal use, which extended to the cabbage-eater's urine, in which infants might be rinsed.^[34] Pliny the Elder listed seven varieties, including Pompeii cabbage, Cumae cabbage and Sabellian cabbage.^[26] According to Pliny, the Pompeii cabbage, which could not stand cold, is "taller, and has a thick stock near the root, but grows thicker between the leaves, these being scantier and narrower, but their tenderness is a valuable quality".^[32] The Pompeii cabbage was also mentioned by Columella in *De Re Rustica*.^[32] Apicius gives several recipes for *cauliculi*, tender cabbage shoots. The Greeks and Romans claimed medicinal usages for their cabbage varieties that included relief from gout, headaches and the symptoms of poisonous mushroom ingestion.^[35] The antipathy towards the vine made it seem that eating cabbage would enable one to avoid drunkenness.^[36] Cabbage continued to figure in the *materia medica* of antiquity as well as at table: in the first century AD Dioscorides mentions two kinds of coleworts with medical uses, the cultivated and the wild,^[11] and his opinions continued to be paraphrased in herbals right through the 17th century

At the end of Antiquity cabbage is mentioned in *De observatione ciborum* ("On the Observance of Foods") of Anthimus, a Greek doctor at the court of Theodoric the Great, and cabbage appears among vegetables directed to be cultivated in the *Capitulare de villis*, composed c. 771-800 that guided the governance of the royal estates of Charlemagne.

In Britain, the Anglo-Saxons cultivated *cawel*.^[37] When round-headed cabbages appeared in 14th-century England they were called *cabaches* and *caboches*, words drawn from Old French and applied at first to refer to the ball of unopened leaves,^[38] the contemporaneous recipe that commences "Take cabbages and quarter them, and seethe them in good broth",^[39] also suggests the tightly headed cabbage.

Manuscript illuminations show the prominence of cabbage in the cuisine of the High Middle Ages,^[21] and cabbage seeds feature among the seed list of purchases for the use of King John II of France when captive in England in 1360,^[40] but cabbages were also a familiar staple of the poor: in the lean year of 1420 the "Bourgeois of Paris" noted that "poor people ate no bread, nothing but cabbages and turnips and such dishes, without any bread or salt".^[41] French naturalist Jean Ruel made what is considered the first explicit mention of head cabbage in his 1536 botanical treatise *De Natura Stirpium*, referring to it as *capucos coles* ("head-coles"),^[42] Sir Anthony Ashley, 1st Baronet, did not disdain to have a cabbage at the foot of his monument in Wimborne St Giles.^[43]

In Istanbul Sultan Selim III penned a tongue-in-cheek ode to cabbage: without cabbage the halva feast was not complete.^[44] Cabbages spread from Europe into Mesopotamia and Egypt as a winter vegetable, and later followed trade routes throughout Asia and the Americas.^[24] The absence of Sanskrit or other ancient Eastern language names for cabbage suggests that it was introduced to South Asia relatively recently.^[6] In India, cabbage was one of several vegetable crops introduced by colonizing traders from Portugal, who established trade routes from the 14th to 17th centuries.^[45] Carl Peter Thunberg reported that cabbage was not yet known in Japan in 1775.^[11]

Many cabbage varieties—including some still commonly grown—were introduced in Germany, France, and the Low Countries.^[6] During the 16th century German gardeners developed the savoy cabbage.^[46] During the 17th and 18th centuries, cabbage was a food staple in such countries as Germany, England, Ireland and Russia, and pickled cabbage was frequently eaten.^[47] Sauerkraut was used by Dutch, Scandinavian and German sailors to prevent scurvy during long ship voyages.^[48]

Jacques Cartier first brought cabbage to the Americas in 1541–42, and it was probably planted by the early English colonists, despite the lack of written evidence of its existence there until the mid-17th century. By the 18th century, it was commonly planted by both colonists and native American Indians.^[6] Cabbage seeds traveled to Australia in 1788 with the First Fleet, and were planted the same year on Norfolk Island. It became a favorite vegetable of Australians by the 1830s and was frequently seen at the Sydney Markets.^[46]



Harvesting cabbage, *Tacuinum Sanitatis*, 15th century.

There are several Guinness Book of World Records entries related to cabbage. These include the heaviest cabbage, at 57.61 kilograms (127.0 lb),^[49] heaviest red cabbage, at 19.05 kilograms (42.0 lb),^[50] longest cabbage roll, at 15.37 meters (50.4 ft),^[51] and the largest cabbage dish, at 925.4 kilograms (2,040 lb).^[52] In 2012, Scott Robb of Palmer, Alaska, broke the world record for heaviest cabbage at 62.71 kilograms (138.25 lb).^[53]

Cultivation



A cabbage field

Cabbage is generally grown for its densely leaved heads, produced during the first year of its biennial cycle. Plants perform best when grown in well-drained soil in a location that receives full sun. Different varieties prefer different soil types, ranging from lighter sand to heavier clay, but all prefer fertile ground with a pH between 6.0 and 6.8.^[54] For optimal growth, there must be adequate levels of nitrogen in the soil, especially during the early head formation stage, and sufficient phosphorus and potassium during the early stages of expansion of the outer leaves.^[55] Temperatures between 4 and 24 °C (39 and 75 °F) prompt the best growth, and extended periods of higher or lower temperatures may result in premature bolting (flowering).^[54]

Flowering induced by periods of low temperatures (a process called vernalization) only occurs if the plant is past the juvenile period. The transition from a juvenile to adult state happens when the stem diameter is about 6 mm (0.24 in). Vernalization allows the plant to grow to an adequate size before flowering. In certain climates, cabbage can be planted at the beginning of the cold period and survive until a later warm period without being induced to flower, a practice that was common in the eastern US.^[56]

Plants are generally started in protected locations early in the growing season before being transplanted outside, although some are seeded directly into the ground from which they will be harvested.^[15] Seedlings typically emerge in about 4–6 days from seeds planted 1.3 cm (0.5 in) deep at a soil temperature between 20 and 30 °C (68 and 86 °F).^[57] Growers normally place plants 30 to 61 cm (12 to 24 in) apart.^[15] Closer spacing reduces the resources available to each plant (especially the amount of light) and increases the time taken to reach maturity.^[58] Some varieties of cabbage have been developed for ornamental use; these are generally called "flowering cabbage". They do not produce heads and feature purple or green outer leaves surrounding an inner grouping of smaller leaves in white, red, or pink.^[15] Early varieties of cabbage take about 70 days from planting to reach maturity, while late varieties take about 120 days.^[59] Cabbages are mature when they are firm and solid to the touch. They are harvested by cutting the stalk just below the bottom leaves with a blade. The outer leaves are trimmed, and any diseased, damaged, or necrotic leaves are removed.^[60] Delays in harvest can result in the head splitting as a result of expansion of the inner leaves and continued stem growth.^[61] Factors that contribute to reduced head weight include: growth in the compacted soils that result from no-till farming practices, drought, waterlogging, insect and disease incidence, and shading and nutrient stress caused by weeds.^[55]



Green and purple cabbages

When being grown for seed, cabbages must be isolated from other *B. oleracea* subspecies, including the wild varieties, by 0.8 to 1.6 km (0.5 to 1 mi) to prevent cross-pollination. Other Brassica species, such as *B. rapa*, *B. juncea*, *B. nigra*, *B. napus* and *Raphanus sativus*, do not readily cross-pollinate.^[62]

Cultivars

There are several cultivar groups of cabbage, each including many cultivars:

- Savoy – Characterized by crimped or curly leaves, mild flavor and tender texture.^[21]
- Spring Greens – Loose-headed, commonly sliced and steamed.^[21]
- Green – Light to dark green, slightly pointed heads.^[21] This is the most commonly grown cultivar.^[15]
- Red – Smooth red leaves, often used for pickling or stewing.^[21]
- White, also called Dutch – Smooth, pale green leaves.^[21]

Some sources only delineate three cultivars: savoyred and white, with spring greens and green cabbage being subsumed into the latter^[63]

Cultivation problems

Due to its high level of nutrient requirements, cabbage is prone to nutrient deficiencies, including boron, calcium, phosphorus and potassium.^[54] There are several physiological disorders that can affect the postharvest appearance of cabbage. Internal tip burn occurs when the margins of inside leaves turn brown, but the outer leaves look normal. Necrotic spot is where there are oval sunken spots a few millimeters across that are often grouped around the midrib. In pepper spot, tiny black spots occur on the areas between the veins, which can increase during storage.^[64]



White cabbage

Fungal diseases include wirestem, which causes weak or dying transplants; Fusarium yellows, which result in stunted and twisted plants with yellow leaves; and blackleg (see Leptosphaeria maculans), which leads to sunken areas on stems and gray-brown spotted leaves.^[65] The fungi Alternaria brassicae and A. brassicicola cause dark leaf spots in affected plants. They are both seedborne and airborne, and typically propagate from spores in infected plant debris left on the soil surface for up to twelve weeks after harvest. Rhizoctonia solani causes the post-emergence disease wirestem, resulting in killed seedlings ("damping-off"), root rot or stunted growth and smaller heads.^[66]

One of the most common bacterial diseases to affect cabbage is black rot, caused by Xanthomonas campestris, which causes chlorotic and necrotic lesions that start at the leaf margins, and wilting of plants. Clubroot, caused by the soilborne slime mold-like organism Plasmodiophora brassicae, results in swollen, club-like roots. Downy mildew, a parasitic disease caused by the oomycete Peronospora parasitica,^[66] produces pale leaves with white, brownish or olive mildew on the lower leaf surfaces; this is often confused with the fungal disease powdery mildew.^[65]



Cabbage moth damage to a savoy cabbage

Pests include root-knot nematodes and cabbage maggots, which produce stunted and wilted plants with yellow leaves; aphids, which induce stunted plants with curled and yellow leaves; harlequin bugs, which cause white and yellow leaves; thrips, which lead to leaves with white-bronze spots; striped flea beetles, which riddle leaves with small holes; and caterpillars, which leave behind large, ragged holes in leaves.^[65] The caterpillar stage of the "small cabbage white butterfly" (Pieris rapae), commonly known in the United States as the "imported cabbage worm", is a major cabbage pest in most countries. The large white butterfly (Pieris brassicae) is prevalent in eastern European countries. The diamondback moth (Plutella xylostella) and the cabbage moth (Mamestra brassicae) thrive in the higher summer temperatures of continental Europe, where they cause considerable damage to cabbage crops.^[67] The cabbage looper (Trichoplusia ni) is infamous in North America for its voracious appetite and for producing frass that contaminates plants.^[68] In India, the diamondback moth has caused losses up to 90 percent in crops that were not treated with insecticide.^[69] Destructive soil insects include the cabbage root fly (Delia radicum) and the cabbage maggot (Hylemya brassicae), whose larvae can burrow into the part of plant consumed by humans.^[67]

Planting near other members of the cabbage family, or where these plants have been placed in previous years, can prompt the spread of pests and disease.^[54] Excessive water and excessive heat can also cause cultivation problems.^[65]

Production

In 2014, global production of cabbages (combined with other brassicas) was 71.8 million tonnes, led by China with 47% of the world total (table). Other major producers were India, Russia, and South Korea.^[70]

Cabbages sold for market are generally smaller, and different varieties are used for those sold immediately upon harvest and those stored before sale. Those used for processing, especially sauerkraut, are larger and have a lower percentage of water.^[16] Both hand and mechanical harvesting are used, with hand-harvesting generally used for cabbages destined for market sales. In commercial-scale operations, hand-harvested cabbages are trimmed, sorted, and packed directly in the field to increase efficiency. Vacuum cooling rapidly refrigerates the vegetable, allowing for earlier shipping and a fresher product. Cabbage can be stored the longest at −1 to 2 °C (30 to 36 °F) with a humidity of 90–100 percent; these conditions will result in up to six months of longevity. When stored under less ideal conditions, cabbage can still last up to four months.^[71]

Culinary use

Cabbage consumption varies widely around the world: Russia has the highest annual per capita consumption at 20 kilograms (44 lb), followed by Belgium at 4.7 kilograms (10 lb), the Netherlands at 4.0 kilograms (8.8 lb), and Spain at 1.9 kilograms (4.2 lb). Americans consume 3.9 kilograms (8.6 lb) annually per capita.^{[35][72]}

Cabbage is prepared and consumed in many ways. The simplest options include eating the vegetable raw or steaming it, though many cuisines pickle, stew, sauté or braise cabbage.^[21] Pickling is one of the most popular ways of preserving cabbage, creating dishes such as sauerkraut and kimchi,^[15] although kimchi is more often made from Chinese cabbage (*B. rapa*).^[21] Savoy cabbages are usually used in salads, while smooth-leaf types are utilized for both market sales and processing.^[16] Bean curd and cabbage is a staple of Chinese cooking,^[73] while the British dish bubble and squeak is made primarily with leftover potato and boiled cabbage and eaten with cold meat.^[74] In Poland, cabbage is one of the main food crops, and it features prominently in Polish cuisine. It is frequently eaten, either cooked or as sauerkraut, as a side dish or as an ingredient in such dishes as bigos (cabbage, sauerkraut, meat, and wild mushrooms, among other ingredients) gołąbki (stuffed cabbage) and pierogi (filled dumplings). Other eastern European countries, such as Hungary and Romania, also have traditional dishes that feature cabbage as a main ingredient.^[75] In India and Ethiopia, cabbage is often included in spicy salads and braises.^[76] In the United States, cabbage is used primarily for the production of coleslaw, followed by market use and sauerkraut production.^[35]

The characteristic flavor of cabbage is caused by glucosinolates, a class of sulfur-containing glucosides. Although found throughout the plant, these compounds are concentrated in the highest quantities in the seeds; lesser quantities are found in young vegetative tissue, and they decrease as the tissue ages.^[77] Cooked cabbage is often criticized for its pungent, unpleasant odor and taste. These develop when cabbage is overcooked and hydrogen sulfide gas is produced.^[78]

Nutrients and phytochemicals

Cabbage is a rich source of vitamin C and vitamin K, containing 44% and 72%, respectively, of the Daily Value (DV) per 100-gram amount (right table of USDA nutrient values).^[79] Cabbage is also a moderate source (10–19% DV) of vitamin

Cabbage production – 2014	
Country	Production (millions of tonnes)
 China	33.4
 India	9.0
 Russia	3.5
 South Korea	2.9
 Ukraine	1.9
 Japan	1.5
World	71.8
Source: FAOSTAT of the United Nations ^[70]	

Cabbage, raw

Nutritional value per 100 g (3.5 oz)	
Energy	103 kJ (25 kcal)
Carbohydrates	5.8 g
Sugars	3.2 g
Dietary fiber	2.5 g
Fat	0.1 g
Protein	1.28 g
Vitamins	
Thiamine (B ₁)	0.061 mg (5%)
Riboflavin (B ₂)	0.040 mg (3%)
Niacin (B ₃)	0.234 mg (2%)
Pantothenic acid (B ₅)	0.212 mg (4%)
Vitamin B ₆	0.124 mg(10%)
Folate (B ₉)	43 µg (11%)
Vitamin C	36.6 mg (44%)
Vitamin K	76 µg (72%)
Minerals	
Calcium	40 mg (4%)
Iron	0.47 mg (4%)
Magnesium	12 mg (3%)
Manganese	0.16 mg (8%)
Phosphorus	26 mg (4%)
Potassium	170 mg (4%)
Sodium	18 mg (1%)
Zinc	0.18 mg (2%)
Other constituents	
Fluoride	1 µg
Link to USDA Database entry	
Units	

B6 and folate, with no other nutrients having significant content per 100-gram serving (table).

µg = micrograms • mg = milligrams
IU = International units

Percentages are roughly approximated
using US recommendations for adults.

Source: USDA Nutrient Database

Basic research on cabbage phytochemicals is ongoing to discern if certain cabbage compounds may affect health or have anti-disease effects. Such compounds include sulforaphane and other glucosinolates which may stimulate the production of detoxifying enzymes during metabolism.^[80] Studies suggest that cruciferous vegetables, including cabbage, may have protective effects against colon cancer.^[81] Cabbage is a source of indole-3-carbinol, a chemical under basic research for its possible properties.^[82]

Herbal medicine

In addition to its usual purpose as an edible vegetable, cabbage has been used historically as a medicinal herb for a variety of purported health benefits. For example, the Ancient Greeks recommended consuming the vegetable as a laxative,^[42] and used cabbage juice as an antidote for mushroom poisoning^[83] for eye salves, and for liniments used to help bruises heal.^[84] In *De Agri Cultura* (*On Agriculture*), Cato the Elder suggested that women could prevent diseases by bathing in urine obtained from those who had frequently eaten cabbage.^[42] The ancient Roman nobleman Pliny the Elder described both culinary and medicinal properties of the vegetable, recommending it for drunkenness—both preventatively to counter the effects of alcohol and to cure hangovers.^[85] Similarly, the Ancient Egyptians ate cooked cabbage at the beginning of meals to reduce the intoxicating effects of wine.^[86] This traditional usage persisted in European literature until the mid-20th century.^[87]

The cooling properties of the leaves were used in Britain as a treatment for trench foot in World War I, and as compresses for ulcers and breast abscesses. Accumulated scientific evidence corroborates that cabbage leaf treatment can reduce the pain and hardness of engorged breasts, and increase the duration of breast feeding.^[88] Other medicinal uses recorded in European folk medicine include treatments for rheumatism, sore throat, hoarseness, colic, and melancholy.^[87] In the United States, cabbage has been used as a hangover cure, to treat abscesses, to prevent sunstroke, or to cool body parts affected by fevers. The leaves have also been used to soothe sore feet and, when tied around a child's neck, to relieve croup. Both mashed cabbage and cabbage juice have been used in poultices to remove boils and treat warts, pneumonia, appendicitis, and ulcers.^[87]

Disadvantages

Bloating

Excessive consumption of cabbage may lead to increased intestinal gas which causes bloating and flatulence due to the trisaccharide raffinose, which the human small intestine cannot digest.^[89]

Food-borne illness

Cabbage has been linked to outbreaks of some food-borne illnesses, including Listeria monocytogenes^[90] and Clostridium botulinum. The latter toxin has been traced to pre-made, packaged coleslaw mixes, while the spores were found on whole cabbages that were otherwise acceptable in appearance. Shigella species are able to survive in shredded cabbage.^[91] Two outbreaks of E. coli in the United States have been linked to cabbage consumption. Biological risk assessments have concluded that there is the potential for further outbreaks linked to uncooked cabbage, due to contamination at many stages of the growing, harvesting and packaging processes. Contaminants from water, humans, animals and soil have the potential to be transferred to cabbage, and from there to the end consumer.^[92]

Goiter and iodine intake

Cabbage and other cruciferous vegetables contain small amounts of thiocyanate, a compound associated with goiter formation when iodine intake is deficient.^[93]

See also

- List of cabbage dishes

References

1. "Classification for species *Brassica oleracea* L." (<http://plants.usda.gov/java/ClassificationServlet?source=profile&symbol=BROL&display=31>) *PLANTS database*. United States Department of Agriculture Retrieved 2012-08-10.
2. Delahaut, K. A.; Newenhouse, A. C (1997). "Growing broccoli, cauliflower cabbage and other cole crops in Wisconsin" (<http://learningstore.uwex.edu/assets/pdfs/A3684.PDF>) (PDF). University of Wisconsin. p. 1 Retrieved 2012-08-12
3. "*Brassica oleracea* L. – Cabbage" (<http://plants.usda.gov/java/profile?symbol=BROL>) United States Department of Agriculture. Retrieved 2012-08-10.
4. Small, Ernst (2009). *Top 100 Food Plants* (https://books.google.com/books?id=nyWY_YkV7qAC&pg=PA127). NRC Research Press. p. 127. ISBN 978-0-660-19858-3
5. "*Brassica oleracea*" (<https://npgsweb.ars-grin.gov/gringlobal/taxonomydetail.aspx?7668>) *Germplasm Resources Information Network (GRIN)*. Agricultural Research Service (ARS), United States Department of Agriculture (USDA). Retrieved 2012-08-12
6. "Of Cabbages and Celts" (<http://aggie-horticulture.tamu.edu/archives/parsons/publications/vegetabletravelers/cabbage.html>). *Aggie Horticulture*. Texas A&M University. Retrieved 2013-10-19.
7. Schneider, Elizabeth (2001). *Vegetables from Amaranth to Zucchini: The Essential Reference* (<https://books.google.com/books?id=cWYWGlcmvO0C&pg=PA196>). HarperCollins. pp. 195–196. ISBN 0-688-15260-0
8. Morris, Charles (1915). *Winston's Cumulative Encyclopedia: A Comprehensive Reference Book* (<https://books.google.com/books?id=ygkoAAAAAAJ&pg=PT337>) 2. J. C. Winston. p. 337.
9. Winer, Lise (2009). *Dictionary of the English/Creole of Trinidad & Tobago: On Historical Principles* (https://books.google.com/books?id=_n82hsbDJBMC&pg=PA150). McGill-Queen's Press. p. 150. ISBN 978-0-7735-3406-3
10. Katz and Weaver, p. 279
11. Sturtevant, Edward Lewis (1919). *Sturtevant's Notes on Edible Plants* (<https://www.biodiversitylibrary.org/page/20118475>). J.B. Lyon. pp. 115, 117.
12. Chantrell, Glynnis, ed. (2002). *The Oxford Dictionary of Word Histories* Oxford University Press. p. 76. ISBN 978-0-19-863121-7.
13. Green, Jonathon (2006). *Cassell's Dictionary of Slang* (<https://books.google.com/books?id=5GpLcC4a5fAC&pg=PA230>). Sterling Publishing Company pp. 230–231. ISBN 978-0-304-36636-1
14. Dixon, p. 19
15. "Cabbage" (<http://urbanext.illinois.edu/veggies/cabbage.cfm>) University of Illinois Extension Retrieved 2012-08-10.
16. Katz and Weaver, p. 280
17. Ordas and Cartea, p. 128
18. Ordas and Cartea, p. 135
19. "Cabbage" (<https://web.archive.org/web/20131018021027/http://www.gmo-compass.org/eng/database/plants/46.cabbage.html>). *GMO Food Database* GMO Compass. Archived from the original (<http://www.gmo-compass.org/eng/database/plants/46.cabbage.html>) on 2013-10-18 Retrieved 2013-10-19.
20. A brief historical sketch is in Maguelonne Tussaint-Samat, *A History of Food*, 2nd ed. 2009, pp 622f.
21. Ingram, Christine (2000). *The Cook's Guide to Vegetables*. Hermes House. pp. 64–66. ISBN 978-1-84038-842-8
22. Dixon, p. 2
23. Chen, S.; Nelson, M. N.; Chèvre, A.- M.; Jenczewski, E.; Li, Z.; Mason, A.; Meng, J.; Plummer, A.; Pradhan, A.; Siddique, K. H. M.; Snowdon, R. J.; Yan, G.; Zhou, W.; Cowling W. A. (2011). "Trigenomic bridges for *Brassica* improvement". *Critical Reviews in Plant Science* **30** (6): 524–547. doi:10.1080/07352689.2011.615700 (<https://doi.org/10.1080/07352689.2011.615700>)
24. Katz and Weaver, p. 284
25. "Cabbage plants" are mentioned by A. Leo Oppenheim *Ancient Mesopotamia: Portrait of a Dead Civilization* rev. ed. 1977, p 313.

26. *Encyclopedia of Cultivated Plants: From Acacia to Zinnia*. ABC-CLIO. 2013. p. 169. ISBN 978-1-59884-775-8
27. Janick, Jules (2011). *Plant Breeding Reviews* (<https://books.google.com/books?id=XAJXJU8Ud8sC&pg=PT51>) 35. John Wiley & Sons. p. 51. ISBN 978-1-118-10049-3
28. *Selected Papyri*, 186, noted in Alan K. Bowman, *Egypt After the Pharaohs* p 151.
29. Dalby, Andrew (2013). *Food in the Ancient World from A to Z* Routledge. p. 67. ISBN 978-1-135-95422-2
30. *Pliny's Natural History*, 20. 78-83.
31. Toussaint-Samat, p. 622.
32. *The Natural History of Pompeii* Cambridge University Press. 2002. p. 94 ISBN 978-0-521-80054-9
33. Toussaint-Samat, p. 623.
34. Cato, *De agricultura*, CLVI, CLVII; the passages are paraphrased by Pliny the Elder
35. Boriss, Hayley; Kreith, Marcia (February 2006). "Commodity Profile: Cabbage" (http://aic.ucdavis.edu/profiles/Cabbage_2006.pdf) (PDF). University of California – Davis Retrieved 2012-08-21
36. Toussaint-Samat, p. 622.
37. Joseph Bosworth. ed. *An Anglo-Saxon Dictionary*, s.v. "cawel" gives parallels: Scottish *kail*, *kale*; Frisian *koal*, *koet*, Dutch *kool* (fem.); German *kohl* (masc.), etc. but also Welsh *cawl*; Cornish *caul* (masc.); etc. Compare Latin *caulis*.
38. *OED*: "cabbage".
39. Jeffrey L. Forgeng, Will McLean, *Daily Life in Chaucer's England* 2009, p 298.
40. Charles Parrain, "The evolution of agricultural technique" in M. M. Postan, ed. *The Cambridge Economic History of Europe: I. The Agrarian Life of the Middle Ages* p 166.
41. Extracts from *A Parisian Journal, 1405-1449* translated by Janet Shirley from the anonymous *Journal d'un bourgeois de Paris* (Oxford: Clarendon Press, 1968). (<https://pantherfile.uwm.edu/carlin/www/parisjournal1.html>) Archived (<https://web.archive.org/web/20140222204356/https://pantherfile.uwm.edu/carlin/www/parisjournal1.html>) 2014-02-22 at the Wayback Machine
42. Wright, Clifford A. (2001). *Mediterranean Vegetables: A Cook's ABC of Vegetables and Their Preparation in Spain, France, Italy, Greece, Turkey, the Middle East, and North Africa with More Than 200 Authentic Recipes for the Home Cook* (<https://books.google.com/books?id=tka838efZvkC&pg=PR77>). Harvard Common Press. pp. 77–79. ISBN 978-1-55832-196-0
43. Cooper, Anthony Ashley (1st Earl of Shaftesbury) (1859). *Memoirs, Letters, and Speeches, of Anthony Ashley Cooper: With Other Papers* (<https://books.google.com/books?id=DdxUAAAacAAJ&pg=PR3>). John Murray p. 3.
44. A translation is in Mary Isin, *Sherbet and Spice: The Complete Story of Turkish Sweets and Desserts* 2013, p. 146.
45. Dabholkar, A. R. (2006). *General Plant Breeding* (<https://books.google.com/books?id=oxooQMxeYhcC&pg=PR135>). Concept Publishing. p. 135. ISBN 978-81-8069-242-0
46. "Cabbage" (https://web.archive.org/web/20120808031331/http://www.freshforkids.com.au/veg_pages/cabbage/cabbage.html). Sydney Markets, Ltd. Archived from the original (http://www.freshforkids.com.au/veg_pages/cabbage/cabbage.html) on 2012-08-08 Retrieved 2012-08-10.
47. Tannahill, pp. 289–291
48. Nolte, Kurt. "Green Cabbage" (<http://cals.arizona.edu/fps/sites/cals.arizona.edu/fps/files/cotw/cabbage.pdf>) (PDF). University of Arizona Retrieved 2012-08-14.
49. "Heaviest cabbage" (<http://www.guinnessworldrecords.com/worldrecords/1000/heaviest-cabbage>) Guinness World Records. Retrieved 2012-08-09.
50. "Heaviest red cabbage" (<http://www.guinnessworldrecords.com/worldrecords/11000/heaviest-red-cabbage>) Guinness World Records. Retrieved 2012-08-09.
51. "Longest cabbage roll" (<http://www.guinnessworldrecords.com/worldrecords/5000/longest-cabbage-roll>) Guinness World Records. Retrieved 2012-08-09.
52. "Largest cabbage dish" (<http://www.guinnessworldrecords.com/worldrecords/5000/largest-cabbage-dish>) Guinness World Records. Retrieved 2012-08-09.
53. "Alaska Man Rolls Record Cabbage Out Of The Patch" (<https://www.npr.org/2012/09/04/160562400/alaska-man-rolls-record-cabbage-out-of-the-patch>) National Public Radio Retrieved 2012-09-04.
54. Bradley et al., pp. 56–57
55. Wien and Murr, p. 533
56. Wien and Murr, pp. 512–515

57. Maynard and Hochmuth, p. 111
58. Wien and Murr, p. 534
59. Maynard and Hochmuth, p. 415
60. Thompson, A. Keith (2003). *Fruit and Vegetables: Harvesting, Handling and Storage* (2nd ed.). Blackwell Publishing. p. 178. [ISBN 978-1-4051-0619-1](#)
61. Wien and Murr, p. 524
62. Katz and Weaver, p. 282
63. Ordas and Cartea, p. 124
64. Becker, Robert F.; Bjorkmann, Thomas. "Nonpathogenic Disorders of Cabbage" (http://vegetablemdonline.ppath.cornell.edu/factsheets/Crucifers_Nonpathogenic.htm) *Vegetable MD Online* Cornell University: Department of Plant Pathology. Retrieved 2013-08-30.
65. Bradley et al., pp. 57–59
66. Keinath, Anthony P; Cubeta, Marc A.; Langston Jr, David B. (2007). "Cabbage diseases: Ecology and control". In Pimentel, David. *Encyclopedia of Pest Management* 2. CRC Press. pp. 56–59. [ISBN 978-1-4200-5361-6](#)
67. Finch, Stan; Collier, Rosemarry H. (2007). "Cruciferous root crop insects: Ecology and control". In Pimentel, David. *Encyclopedia of Pest Management* 2. CRC Press. pp. 131–134. [ISBN 978-1-4200-5361-6](#)
68. Turini TA, Daugovish O, Koike ST, Natwick ET, Ploeg A, Dara SK, Fennimore SA, Joseph S, LeStrange M, Smith R, Subbarao KV, Westerdahl BB. Revised continuously *UC IPM Pest Management Guidelines Cole Crops* UC ANR Publication 3442. Oakland, CA.
69. Janick, p. 195 (<https://books.google.com/books?id=XAjXJU8Ud8sC&pg=PT195>)
70. "Crops/Regions/World list/Production Quantity (pick lists), Cabbages and other brassicas, 2014" (<http://www.fao.org/faostat/en/#data/QC>) UN Food and Agriculture Organization, Corporate Statistical Database (FAOSTAT). 2017. Retrieved 20 May 2017.
71. Katz and Weaver, p. 285
72. "Cabbage" (<https://web.archive.org/web/20120616045430/http://www.fondation-louisbonduelle.org/france/en/know-your-vegetables/nutritional-assets-of-vegetables/chou-vert.html#axzz2g3z7j2gK>) Louis Bonduelle Foundation. Archived from the original (<http://www.fondation-louisbonduelle.org/france/en/know-your-vegetables/nutritional-assets-of-vegetables/chou-vert.html#axzz24EyyXQ7a>) on 2012-06-16 Retrieved 2012-08-22.
73. Tannahill, p. 146
74. Tannahill, p. 277
75. MacVeigh, Jeremy (2008). *International Cuisine* (<https://books.google.com/books?id=g9KcADhDPqkC&pg=PA53>). Cengage Learning. pp. 53–54. [ISBN 978-1-4180-4965-2](#)
76. Marks, Gil (2008). *Olive Trees and Honey: A Treasury of Vegetarian Recipes from Jewish Communities Around the World* (https://books.google.com/books?id=Lb3MVYVp_9sC&pg=PT392) Houghton Mifflin Harcourt. p. 392. [ISBN 978-0-544-18750-4](#)
77. Katz and Weaver, pp. 282–283
78. Corriher, Shirley O. (2000–2001). "Corriher's Compendium of Ingredients and Cooking Problems" (<https://web.archive.org/web/20081217112600/http://units.sla.org/division/dfan/FFT/vol32no1.pdf>) (PDF). *Food for Thought* 32 (1): 6. Archived from the original (<http://units.sla.org/division/dfan/FFT/vol32no1.pdf>) (PDF) on 2008-12-17.
79. "USDA database table for raw cabbage per 100 g" (<http://ndb.nal.usda.gov/ndb/foods/show/2925>) US Department of Agriculture, National Nutrient Database for Standard Reference, version SR-27. 2014 Retrieved 4 December 2014.
80. Dinkova-Kostova AT, Kostov RV (2012). "Glucosinolates and isothiocyanates in health and disease" *Trends Mol Med*. 18 (6): 337–47. doi:10.1016/j.molmed.2012.04.003 (<https://doi.org/10.1016%2Fj.molmed.2012.04.003>) PMID 22578879 (<https://www.ncbi.nlm.nih.gov/pubmed/22578879>).
81. Tse, G; Eslick, G.D. (2014). "Cruciferous vegetables and risk of colorectal neoplasms: a systematic review and meta-analysis". *Nutrition and Cancer*. 66 (1): 128–139. doi:10.1080/01635581.2014.852686 (<https://doi.org/10.1080%2F01635581.2014.852686>) PMID 24341734 (<https://www.ncbi.nlm.nih.gov/pubmed/24341734>).

82. Wu, Y.; Feng, X.; Jin, Y.; Wu, Z.; Hankey, W.; Paisie, C.; Li, L.; Liu, F.; Barsky, S. H.; Zhang, W.; Ganju, R.; Zou, X. (2010). "A novel mechanism of indole-3-carbinol effects on breast carcinogenesis involves induction of Cdc25A degradation". *Cancer Prevention Research* 3 (7): 818–828. doi:10.1158/1940-6207.CAPR-09-0213(<https://doi.org/10.1158/1940-6207.CAPR-09-0213>) PMID 20587702 (<https://www.ncbi.nlm.nih.gov/pubmed/20587702>). Lay summary (<https://www.sciencedaily.com/releases/2010/06/100629131316.htm>)– *Science Daily* (June 30, 2010).
83. Decoteau, Dennis R. (2000). *Vegetable Crops*. Prentice Hall. p. 174. ISBN 978-0-13-956996-8
84. Phillips, Henry (1827). *History of Cultivated Vegetables: Comprising their Botanical, Medicinal, Edible, and Chemical Qualities; Natural History*(<https://books.google.com/books?id=pDkAAAAAJ&pg=PA99>). Henry Colburn. p. 99.
85. Dalby, Andrew; Grainger, Sally (1996). *The Classical Cookbook*(<https://books.google.com/books?id=T7S5iC3pZpOC&pg=PA52>). Getty Publications. p. 52. ISBN 978-0-89236-394-0
86. Janick, p. 51 (<https://books.google.com/books?id=XAJXJU8Ud8sC&pg=PT51>)
87. Hatfield, Gabrielle (2004). *Encyclopedia of Folk Medicine: Old World and New World Editions* (<https://books.google.com/books?id=2GGz6708nqgC&pg=PA60>). ABC-CLIO. pp. 59–60. ISBN 978-1-57607-874-7.
88. Wong, Boh Boi; Koh, Serena; Hegney Desley Gail; He, Hong-Gu (2012). "The effectiveness of cabbage leaf application (treatment) on pain and hardness in breast engorgement and its effect on the duration of breastfeeding" (<https://web.archive.org/web/20131002131620/http://www.joannabriggslibrary.org/jbilibrary/index.php/jbisrir/article/view/58>). *The JBI Database of Systematic Reviews and Implementation Reports* 10 (20): 1185–1213. Archived from the original (<http://www.joannabriggslibrary.org/jbilibrary/index.php/jbisrir/article/view/58>) on 2013-10-02.
89. "The digestive system and gas"(<http://www.webmd.com/digestive-disorders/digestive-diseases-gas>) WebMD. Retrieved 24 June 2013.
90. Davis, J. G.; Kendall, P (April 19, 2013). "Preventing *E. coli* from Garden to Plate"(<http://www.ext.colostate.edu/pub/s/foodnut/09369.html>) Colorado State University Retrieved 2012-08-10.
91. "Chapter IV. Outbreaks Associated with Fresh and Fresh-Cut Produce. Incidence, Growth, and Survival of Pathogens in Fresh and Fresh-Cut Produce"(<https://web.archive.org/web/20121109123857/http://www.fda.gov/Food/ScienceResearch/ResearchAreas/SafePracticesforFoodProcesses/ucm091265.htm>) *Analysis and Evaluation of Preventive Control Measures for the Control and Reduction/Elimination of Microbial Hazards on Fresh and Fresh-Cut Produce*. US Food and Drug Administration. April 12, 2012. Archived from the original (<http://www.fda.gov/Food/ScienceResearch/ResearchAreas/SafePracticesforFoodProcesses/ucm091265.htm>) on November 9, 2012 Retrieved 2012-08-10.
92. "Cabbage Risk Assessment Introduction and Summary"(https://web.archive.org/web/20130921055715/http://www.omafra.gov.on.ca/english/food/inspection/fruitveg/risk_assessment_pdf/cabbage/30ra.pdf) (PDF). Ontario Ministry of Agriculture, Food and Rural Affairs. December 2001. Archived from the original (http://www.omafra.gov.on.ca/english/food/inspection/fruitveg/risk_assessment_pdf/cabbage/30ra.pdf) (PDF) on 2013-09-21 Retrieved 2012-08-19.
93. Vanderpas J (2006). "Nutritional epidemiology and thyroid hormone metabolism"(http://www.iccidd.org/cm_data/2006_Vanderpas_Nutritional_epidemiology_andthyroid_hormone_metabolism_AnnRevNutr.pdf) (PDF). *Annu Rev Nutr.* 26: 293–322. doi:10.1146/annurevnutr.26.010506.103810(<https://doi.org/10.1146/annurevnutr.26.010506.103810>). PMID 16704348 (<https://www.ncbi.nlm.nih.gov/pubmed/16704348>).

Works cited

- Bradley, Fern Marshall; Ellis, Barbara W; Martin, Deborah L., eds. (2009). *The Organic Gardener's Handbook of Natural Pest and Disease Control* Rodale, Inc. ISBN 978-1-60529-677-7.
- Dixon, Geoffrey R. (2007). *Vegetable Brassicas and Related Crucifers* Crop Production Science in Horticulture 14. CAB International. ISBN 978-0-85199-395-9
- Janick, Jules (2011). *Plant Breeding Reviews* 35. John Wiley & Sons. ISBN 978-1-118-10049-3
- Katz, Solomon H.; Weaver, William Woys (2003). *Encyclopedia of Food and Culture* 2. Scribner. ISBN 978-0-684-80565-8.
- Maynard, Donald N.; Hochmuth, George J. (2007) *Knott's Handbook for Vegetable Growers* (5th ed.). Wiley ISBN 978-0-471-73828-2
- Ordas, Amando; Cartea, M. Elena (2008). "Cabbage and Kale". In Prohens, J.; Nuez, J. *Vegetables I: Asteraceae, Brassicaceae, Chenopodiaceae, and Cucurbitaceae* 2. Springer. ISBN 978-0-387-72291-7.
- Tannahill, Reay (1973). *Food in History*. Stein and Day ISBN 978-0-8128-1437-8
- Wien, H. C.; Murr, D. C. E. (1997). "Cauliflower, broccoli, cabbage and brussel sprouts". In Wien, H. C. *The Physiology of Vegetable Crops*. CAB International. ISBN 978-0-85199-146-7.

External links

- *Cabbages and Cauliflowers: How to Grow Them* by James John Howard Gregory – Project Gutenberg.
-

Retrieved from '<https://en.wikipedia.org/w/index.php?title=Cabbage&oldid=834080646>

This page was last edited on 3 April 2018, at 20:45.

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.