


Line–line intersection

In [Euclidean geometry](#), the [intersection](#) of a [line](#) and a line can be the [empty set](#), a [point](#), or a line. Distinguishing these cases and finding the intersection point have use, for example, in [computer graphics](#), [motion planning](#) and [collision detection](#).

In [three-dimensional](#) Euclidean geometry, if two lines are not in the same [plane](#) they are called [skew lines](#) and have no point of intersection. If they are in the same plane there are three possibilities: if they coincide (are not distinct lines) they have an [infinitude](#) of points in common (namely all of the points on either of them); if they are distinct but have the same slope they are said to be [parallel](#) and have no points in common; otherwise they have a single point of intersection.



The intersection of lines.

The distinguishing features of [non-Euclidean geometry](#) are the number and locations of possible intersections between two lines and the number of possible lines with no intersections (parallel lines) with a given line.

Contents

Intersection of two lines

- Given two points on each line
- Given the equations of the lines
- Using homogeneous coordinates

n-line intersection

- Existence of and expression for the intersection
 - In two dimensions
 - In three dimensions
- Nearest point to non-intersecting lines
 - In two dimensions
 - In three dimensions

See also

References

External links

Intersection of two lines

A [necessary condition](#) for two lines to intersect is that they are in the same plane—that is, are not skew lines. Satisfaction of this condition is equivalent to the [tetrahedron](#) with vertices at two of the points on one line and two of the points on the other line being [degenerate](#) in the sense of having [zero volume](#). For the algebraic form of this condition, see [Skew lines § Testing for skewness](#)

Given two points on each line

First we consider the intersection of two lines L_1 and L_2 in 2-dimensional space, with line L_1 being defined by two distinct points (x_1, y_1) and (x_2, y_2) , and line L_2 being defined by two distinct points (x_3, y_3) and (x_4, y_4) .^[1]

The intersection P of line L_1 and L_2 can be defined using determinants.

$$P_x = \frac{\begin{vmatrix} x_1 & y_1 \\ x_2 & y_2 \end{vmatrix} \begin{vmatrix} x_1 & 1 \\ x_2 & 1 \end{vmatrix} - \begin{vmatrix} x_3 & y_3 \\ x_4 & y_4 \end{vmatrix} \begin{vmatrix} x_3 & 1 \\ x_4 & 1 \end{vmatrix}}{\begin{vmatrix} x_1 & 1 \\ x_2 & 1 \end{vmatrix} \begin{vmatrix} y_1 & 1 \\ y_2 & 1 \end{vmatrix} - \begin{vmatrix} x_3 & 1 \\ x_4 & 1 \end{vmatrix} \begin{vmatrix} y_3 & 1 \\ y_4 & 1 \end{vmatrix}}, \quad P_y = \frac{\begin{vmatrix} x_1 & y_1 \\ x_2 & y_2 \end{vmatrix} \begin{vmatrix} y_1 & 1 \\ y_2 & 1 \end{vmatrix} - \begin{vmatrix} x_3 & y_3 \\ x_4 & y_4 \end{vmatrix} \begin{vmatrix} y_3 & 1 \\ y_4 & 1 \end{vmatrix}}{\begin{vmatrix} x_1 & 1 \\ x_2 & 1 \end{vmatrix} \begin{vmatrix} y_1 & 1 \\ y_2 & 1 \end{vmatrix} - \begin{vmatrix} x_3 & 1 \\ x_4 & 1 \end{vmatrix} \begin{vmatrix} y_3 & 1 \\ y_4 & 1 \end{vmatrix}}$$

The determinants can be written out as:

$$(P_x, P_y) = \left(\frac{(x_1 y_2 - y_1 x_2)(x_3 - x_4) - (x_1 - x_2)(x_3 y_4 - y_3 x_4)}{(x_1 - x_2)(y_3 - y_4) - (y_1 - y_2)(x_3 - x_4)}, \frac{(x_1 y_2 - y_1 x_2)(y_3 - y_4) - (y_1 - y_2)(x_3 y_4 - y_3 x_4)}{(x_1 - x_2)(y_3 - y_4) - (y_1 - y_2)(x_3 - x_4)} \right)$$

Note that the intersection point is for the infinitely long lines defined by the points, rather than the line segments between the points, and can produce an intersection point beyond the lengths of the line segments. If (rather than solving for the point in a single step), the solution in terms of first degree Bézier parameters is first found, then this intermediate result can be checked for $0.0 \leq t \leq 1.0$ and $0.0 \leq u \leq 1.0$ (where t and u are the driving variables).

When the two lines are parallel or coincident the denominator is zero:

$$(x_1 - x_2)(y_3 - y_4) - (y_1 - y_2)(x_3 - x_4) = 0.$$

If the lines are almost parallel, then a computer solution might encounter numeric problems implementing the solution described above: the recognition of this condition might require an approximate test in a practical application. An alternate approach might be to rotate the line segments so that one of them is horizontal, whence the solution of the rotated parametric form of the second line is easily obtained. Careful discussion of the special cases is required (parallel lines/coincident lines, overlapping/non-overlapping intervals).

Given the equations of the lines

The x and y coordinates of the point of intersection of two non-vertical lines can easily be found using the following substitutions and rearrangements.

Suppose that two lines have the equations $y = ax + c$ and $y = bx + d$ where a and b are the slopes (gradients) of the lines and where c and d are the y -intercepts of the lines. At the point where the two lines intersect (if they do), both y coordinates will be the same, hence the following equality:

$$ax + c = bx + d.$$

We can rearrange this expression in order to extract the value of x ,

$$ax - bx = d - c,$$

and so,

$$x = \frac{d - c}{a - b}.$$

To find the y coordinate, all we need to do is substitute the value of x into either one of the two line equations, for example, into the first:

$$y = a \frac{d - c}{a - b} + c.$$

Hence, the point of intersection is

$$P \left(\frac{d - c}{a - b}, a \frac{d - c}{a - b} + c \right) = P \left(\frac{d - c}{a - b}, \frac{ad - bc}{a - b} \right).$$

Note if $a = b$ then the two lines are parallel. If $c \neq d$ as well, the lines are different and there is no intersection, otherwise the two lines are identical.

Using homogeneous coordinates

By using homogeneous coordinates the intersection point of two implicitly defined lines can be determined quite easily. In 2D, every point can be defined as a projection of a 3D point, given as the ordered triple (x, y, w) . The mapping from 3D to 2D coordinates is $(x', y') = (x/w, y/w)$. We can convert 2D points to homogeneous coordinates by defining them as $(x, y, 1)$.

Assume that we want to find intersection of two infinite lines in 2-dimensional space, defined as $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$. We can represent these two lines in line coordinates as $U_1 = (a_1, b_1, c_1)$ and $U_2 = (a_2, b_2, c_2)$.

The intersection P' of two lines is then simply given by^[2]

$$P' = (a_p, b_p, c_p) = U_1 \times U_2 = (b_1c_2 - b_2c_1, a_2c_1 - a_1c_2, a_1b_2 - a_2b_1)$$

If $c_p = 0$ the lines do not intersect.

n -line intersection

Existence of and expression for the intersection

In two dimensions

In two dimensions, more than two lines almost certainly do not intersect at a single point. To determine if they do and, if so, to find the intersection point, write the i -th equation ($i = 1, \dots, n$) as $(a_{i1} \ a_{i2})(x \ y)^T = b_i$, and stack these equations into matrix form as

$$Aw = b,$$

where the i -th row of the $n \times 2$ matrix A is (a_{i1}, a_{i2}) , w is the 2×1 vector $(x, y)^T$, and the i -th element of the column vector b is b_i . If A has independent columns, its rank is 2. Then if and only if the rank of the augmented matrix $[A \mid b]$ is also 2, there exists a solution of the matrix equation and thus an intersection point of the n lines. The intersection point, if it exists, is given by

$$w = A^g b = (A^T A)^{-1} A^T b,$$

where A^g is the Moore-Penrose generalized inverse of A (which has the form shown because A has full column rank). Alternatively, the solution can be found by jointly solving any two independent equations. But if the rank of A is only 1, then if the rank of the augmented matrix is 2 there is no solution but if its rank is 1 then all of the lines coincide with each other

In three dimensions

The above approach can be readily extended to three dimensions. In three or more dimensions, even two lines almost certainly do not intersect; pairs of non-parallel lines that do not intersect are called skew lines. But if an intersection does exist it can be found, as follows.

In three dimensions a line is represented by the intersection of two planes, each of which has an equation of the form $(a_{i1} \ a_{i2} \ a_{i3})(x \ y \ z)^T = b_i$. Thus a set of n lines can be represented by $2n$ equations in the 3-dimensional coordinate vector $w = (x, y, z)^T$:

$$Aw = b$$

where now A is $2n \times 3$ and b is $2n \times 1$. As before there is a unique intersection point if and only if A has full column rank and the augmented matrix $[A \mid b]$ does not, and the unique intersection if it exists is given by

$$w = (A^T A)^{-1} A^T b.$$

Nearest point to non-intersecting lines

In two or more dimensions, we can usually find a point that is mutually closest to two or more lines in least-squares sense.

In two dimensions

In the two-dimensional case, first, represent line i as a point, p_i , on the line and a unit normal vector, $\hat{n}_i$, perpendicular to that line. That is, if x_1 and x_2 are points on line 1, then let $p_1 = x_1$ and let

$$\hat{n}_1 := \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} (x_2 - x_1) / \|x_2 - x_1\|$$

which is the unit vector along the line, rotated by 90 degrees.

Note that the distance from a point x to the line $(p, \hat{n})$ is given by

$$d(x, (p, n)) = \|(x - p) \cdot \hat{n}\| = \|(x - p)^T \hat{n}\| = \sqrt{(x - p)^T \hat{n} \hat{n}^T (x - p)}.$$

And so the squared distance from a point x , to a line is

$$d(x, (p, n))^2 = (x - p)^T (\hat{n} \hat{n}^T) (x - p).$$

The sum of squared distances to many lines is the cost function:

$$E(x) = \sum_i (x - p_i)^T (\hat{n}_i \hat{n}_i^T) (x - p_i).$$

This can be rearranged:

$$\begin{aligned}
E(x) &= \sum_i x^\top \hat{n}_i \hat{n}_i^\top x - x^\top \hat{n}_i \hat{n}_i^\top p_i - p_i^\top \hat{n}_i \hat{n}_i^\top x + p_i^\top \hat{n}_i \hat{n}_i^\top p_i \\
&= x^\top \left(\sum_i \hat{n}_i \hat{n}_i^\top \right) x - 2x^\top \left(\sum_i \hat{n}_i \hat{n}_i^\top p_i \right) + \sum_i p_i^\top \hat{n}_i \hat{n}_i^\top p_i.
\end{aligned}$$

To find the minimum, we differentiate with respect to x and set the result equal to the zero vector:

$$\frac{\partial E(x)}{\partial x} = 0 = 2 \left(\sum_i \hat{n}_i \hat{n}_i^\top \right) x - 2 \left(\sum_i \hat{n}_i \hat{n}_i^\top p_i \right)$$

so

$$\left(\sum_i \hat{n}_i \hat{n}_i^\top \right) x = \sum_i \hat{n}_i \hat{n}_i^\top p_i$$

and so

$$x = \left(\sum_i \hat{n}_i \hat{n}_i^\top \right)^{-1} \left(\sum_i \hat{n}_i \hat{n}_i^\top p_i \right).$$

In three dimensions

While $\hat{n}_i$ is not well-defined in more than two dimensions, this can be generalized to any number of dimensions by noting that $\hat{n}_i \hat{n}_i^\top$ is simply the (symmetric) matrix with all eigenvalues unity except for a zero eigenvalue in the direction along the line providing a seminorm on the distance between p_i and another point giving the distance to the line. In any number of dimensions, if $\hat{v}_i$ is a unit vector *along* the i -th line, then

$$\hat{n}_i \hat{n}_i^\top \text{ becomes } I - \hat{v}_i \hat{v}_i^\top$$

where I is the identity matrix, and so

$$x = \left(\sum_i I - \hat{v}_i \hat{v}_i^\top \right)^{-1} \left(\sum_i (I - \hat{v}_i \hat{v}_i^\top) p_i \right).$$

See also

- Line segment intersection
- Line intersection in projective space
- Distance from a point to a line
- Parallel postulate
- Intersection (Euclidean geometry) §Two line segments

References

1. "Weisstein, Eric W. "Line-Line Intersection." From MathWorld" (<http://mathworld.wolfram.com/Line-LineIntersection.html>). *A Wolfram Web Resource*. Retrieved 2008-01-10.
2. "Homogeneous coordinates"(<http://robotics.stanford.edu/~birch/projective/node4.html>)*robotics.stanford.edu* Retrieved 2015-08-18.

External links

- Distance between Lines and Segments with their Closest Point of Approach applicable to two, three, or more dimensions.
-

Retrieved from 'https://en.wikipedia.org/w/index.php?title=Line-line_intersection&oldid=815048738

This page was last edited on 12 December 2017, at 13:04.

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.